

07/733,896

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show 8 Numbers](#)[Edit 8 Numbers](#)[Preferences](#)[Cases](#)**Search Results -**

Term	Documents
IMMUNO.DWPI,EPAB,JPAB,USPT.	16371
IMMUNOES	0
IMMUNOS	0
IMMUNOE.DWPI,EPAB,JPAB,USPT.	3
POLYMERASE.DWPI,EPAB,JPAB,USPT.	31976
POLYMERASES.DWPI,EPAB,JPAB,USPT.	4405
CHAIN.DWPI,EPAB,JPAB,USPT.	647763
CHAINS.DWPI,EPAB,JPAB,USPT.	161455
REACTION.DWPI,EPAB,JPAB,USPT.	1188147
REACTIONS.DWPI,EPAB,JPAB,USPT.	259010
TRANSCRIPTASE\$1	0
(IMMUNO SAME POLYMERASE CHAIN REACTION SAME TRANSCRIPTASE\$1).USPT,JPAB,EPAB,DWPI.	0

[There are more results than shown above. Click here to view the entire set.](#)

US Patents Full-Text Database

US Pre-Grant Publication Full-Text Database

JPO Abstracts Database**EPO Abstracts Database****Derwent World Patents Index**

Database: IBM Technical Disclosure Bulletins

Search:

L5

[Refine Search](#)[Recall Text](#)[Clear](#)**Search History**DATE: Wednesday, February 13, 2002 [Printable Copy](#) [Create Case](#)

Set Name Query

side by side

Hit Count Set Name

result set

DB=USPT,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

<u>L5</u>	immuno same polymerase chain reaction same trancriptase\$1	0	<u>L5</u>
<u>L4</u>	immuno polymerase chain reaction\$1 same electrophoresis same fluorescen\$2 same transcriptase\$1	0	<u>L4</u>
<u>L3</u>	L2 and transcriptase\$1	0	<u>L3</u>
<u>L2</u>	L1 and fluorescen\$2	1	<u>L2</u>
<u>L1</u>	immuno near5 polymerase chain reaction near5 electrophoresis	1	<u>L1</u>

END OF SEARCH HISTORY

End of Result Set

☐ Generate Collection

L8: Entry 1 of 1

File: USPT

Jun 3, 1997

US-PAT-NO: 5635617

DOCUMENT-IDENTIFIER: US 5635617 A

TITLE: Methods and compositions comprising the agfA gene for detection of Salmonella

DATE-ISSUED: June 3, 1997

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Doran; James L.	Brentwood Bay			CAX
Kay; William W.	Victoria			CAX
Collinson; S. Karen	Brentwood Bay			CAX
Clouthier; Sharon C.	Naniamo			CAX

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE	CODE
University of Victoria Innovation & Development Corp.	Victoria			CAX		03

APPL-NO: 8/ 233788 [PALM]

DATE FILED: April 26, 1994

PARENT-CASE:

CROSS-REFERENCE TO RELATED APPLICATIONS This application is a continuation-in-part of U.S. patent application Ser. No. 08/054,452, filed Apr. 26, 1993, abandoned.

INT-CL: [6] C07 H 21/02, C07 H 21/04

US-CL-ISSUED: 536/23.7; 536/23.1

US-CL-CURRENT: 536/23.7; 536/23.1

FIELD-OF-SEARCH: 536/23.7, 536/23.2, 536/24.33, 536/23.1

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

☐ Search Selected☐ Search ALL

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<input type="checkbox"/> 4443549	April 1984	Sadowski	436/548
<input type="checkbox"/> 4689295	August 1987	Taber et al.	435/6
<input type="checkbox"/> 5043264	August 1991	Jikuya et al.	435/6
<input type="checkbox"/> 5147778	September 1992	Nietupski et al.	435/6

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
114 668 A2	August 1984	EPX	
355 989 A2	February 1990	EPX	
383 509 A2	August 1990	EPX	
2234587	February 1991	GBX	
WO86/00993	February 1986	WOX	
WO86/01805	March 1986	WOX	
WO86/04352	July 1986	WOX	
WO88/03957	June 1988	WOX	
WO89/01162	February 1989	WOX	
WO89/05359	June 1989	WOX	
WO90/00624	January 1990	WOX	
WO91/19003	December 1991	WOX	
WO92/01056	January 1992	WOX	
WO92/06197	April 1992	WOX	
WO92/06198	April 1992	WOX	
WO93/04202	March 1993	WOX	
WO93/20231	October 1993	WOX	

OTHER PUBLICATIONS

Doran et al J. Clin. Microbiol 31: 2263-2273 (1993).

Collinson et al., "Aggregative Fimbriae of Salmonella enteritidis," in Abstracts of the 22nd Annual Western Branch Meeting of the Canadian Society for Microbiology, Abstract #I-10, Nov. 14-16, Vancouver B.C., 1991.

Clouthier et al., "Characterization of three fimbrial genes, sefA, B, and C of Salmonella enteritidis," in Abstracts of the 22nd Annual Western Branch Meeting of the Canadian Society for Microbiology, Abstract #I-9, Nov. 14-16, Vancouver B.C., 1991.

Muller et al., "Type 1 Fimbriae of Salmonella enteritidis," Journal of Bacteriology 173(15):4765-4772, 1991.

Collinson et al., "Purification and Characterization of Thin, Aggregative Fimbriae from Salmonella enteritidis," Journal of Bacteriology 173(15): 4773-4781, 1991.

Collinson et al., "Thin, Aggregative Fimbriae Mediate Binding of Salmonella enteritidis to Fibronectin," Journal of Bacteriology 175(1): 12-18, 1993.

Collinson, S. K., L. Emody, P. Doig, K.-H. Muller, T. Trust and W. Kay. 1992. "Thin, Aggregative Fimbriae Mediate Binding of Salmonella enteritidis to Fibronectin." Abstracts of the 92 General Meeting of the American Society for Microbiology, May 26-30, p. 27, B-401, New Orleans, LA.

Muller et al., "Fimbriation Genes of Salmonella enteritidis," Journal of Bacteriology 171(9): 4648-4654, 1989.

Feutrier et al., "Cloning and Expression of a Salmonella enteritidis Fimbrin Gene in Escherichia coli," Journal of Bacteriology 170(9): 4216-4222, 1988.

Feutrier et al., "Purification and Characterization of Fimbriae from Salmonella enteritidis," Journal of Bacteriology 168(1): 221-227, 1986.

Feutrier et al., "Novel Type 1 Fimbriae of Salmonella Enteritidis," in David L. Larke et al. (eds.), Protein-Carbohydrate Interactions in Biological Systems, Academic Press, 1986, pp. 103-107.

Thorns et al., "Detection of a Novel fimbrial Structure on the Surface of Salmonella enteritidis by Using a Monoclonal Antibody," Journal of Clinical Microbiology 28(11): 2409-2414, 1990.

Thorns et al., "Characterisation of monoclonal antibodies against a fimbrial structure of Salmonella enteritidis and certain other serogroup D salmonellae and their application as serotyping reagents," Research in Veterinary Science 53: 300-308, 1992.

Central Veterinary Laboratory, "A New Latex Agglutination Test Kit for the Identification OF Salmonella Enteritidis," SEFEX Product Advertisement.

Widenhorn et al., "Genetic Regulation of the Tricarboxylate Transport Operon (tctI) of Salmonella typhimurium," Journal of Bacteriology 171(8): 4436-4441, 1989.

Widenhorn et al., "Expression of the Divergent Tricarboxylate Transport Operon (tctI)

of *Salmonella typhimurium*," *Journal of Bacteriology* 170(7): 3223-3227, 1988.

Widenhorn et al., "Cloning and Properties of the *Salmonella typhimurium* tricarboxylate Transport Operon in *Escherichia coli*," *Journal of Bacteriology* 170(2): 883-888, 1988.

Sweet et al., "Tricarboxylate-binding Proteins of *Salmonella typhimurium*," *Journal of Biological Chemistry* 259(3): 1586-1592, 1984.

Somers and kay, "Genetic Fine Structure of the Tricarboxylate transport (tct) Locus of *Salmonella typhimurium*," *Mol. Gen. Genet.* 190: 20-26, 1983.

Sweet et al., "Purification and properties of a citrate-binding transport component, the C protien of *Salmonella typhimurium*," *Canadian Journal of Biochemistry* 57: 710-715, 1979.

Andrews, "A Review Of Culture Methods And Their Relation To Rapid Methods For The Detection Of *Salmonella* In Foods," *Food Technology* 39(3): 77-82, 1985.

Rubin, "Nucleic Acid Probes for the Identification of *Salmonella*," in Alberto J.L. Macario et al. (eds.), *Gene Probes for Bacteria*, Academic Press, 1990, pp. 323-352.

Blackburn, "Rapid and alternative methods for the detection of salmonellas in foods," *Journal of Applied Bacteriology* 75: 199-214, 1993.

Mattingly et al., "Use of Monoclonal Antibodies For The Detection of *Salmonella* In Foods," *Food technology* 39(3): 90-94, 1985.

Cox, "Salmonella Methodology Update," *Poultry Science* 67: 921-927, 1988.

Swaminathan et al., "Enzyme Immunoassay For *Salmonella*: One-Day Testing Is Now A Reality," *Food Technology* 39(3): 83-89, 1985.

Fitts, "Development Of A DNA-DNA Hybridization Test For The Presence of *Salmonella* In Foods," *Food Technology* 39(3): 94-102, 1985.

Flowers, "Comparison of Rapid *Salmonella* Screening Methods And The Conventional Culture Method," *Food Technology* 39(3): 103-108, 1985.

Hartman and Minnich, "Automation for Rapid Identification of *Salmonellae* in Foods," *Journal of Food Protection* 44(5): 385-393, 1981.

Thomason, "Current Status of Immunofluorescent Methodology for *Salmonellae*," *Journal of Food Protection* 44(5) 381-384, 1981.

Tompkins et al, "Cloned, Random Chromosomal Sequences as Probes to Identify *Salmonella* Species," *Journal of Infectious Diseases* 154(1): 156-162, 1986.

Fitts et al., "DNA-DNA Hybridization Assay for Detection of *Salmonella* spp. in Foods," *Applied and Environmental Microbiology* 46: 1146-1151, 1983.

Ibrahim et al., "Rapid Detection of *Salmonellae* by Immunoassays with Titanous Hydroxide as the Solid Phase," *Applied and Environmental Microbiology* 50(3): 670-675, 1985.

Rahn et al., "Amplification of an *invA* gene sequence of *Salmonella typhimurium* by polymerase chain reaction as a specific method of detection of *Salmonella*," *Molecular and Cellular Probes* 6: 271-279, 1992.

Chan et al., "Comparative Study of Colorimetric DNA Hybridization Method and Conventional Culture Procedure for Detection of *Salmonella* in Foods," *Journal Assoc. Off. Anal. Chem.* 73(3): 419-424, 1990.

Flowers et al., "DNA Hybridization Assay for Detection of *Salmonella* in Foods: Collaborative Study," *J. Assoc. Off. Anal. Chem.* 70(3): 521-529, 1987.

Aabo et al., "Salmonella identification by the polymerase chain reaction," *Molecular and Cellular Probes* 7: 171-178, 1993.

Spierings et al., "Characterization of the *Salmonella typhimurium* *phoE* gene and development of *Salmonella*-specific DNA probes," *Gene* 122: 45-52, 1992.

Widjojoatmodjo et al., "The Magnetic Immuno Polymerase Chain Reaction Assay for Direct Detection of *Salmonellae* in Fecal Samples," *Journal of Clinical Microbiology* 30(12): 3195-3199, 1992.

Cano et al., "Detection of salmonellas by DNA hybridization with a fluorescent alkaline phosphatase substrate," *Journal of Applied Bacteriology* 72: 393-399, 1992.

Knight et al., "Direct Detection of *Salmonella* spp. in Estuaries by Using a DNA Probe," *Applied and Environmental Microbiology* 56(4): 1059-1066, 1990.

Gopo et al., "Development of a *Salmonella*-specific biotinylated DNA probe for rapid routine identification of *Salmonella*," *Molecular and Cellular Probes* 2: 271-279, 1988.

Olsen et al., "Isolation of a *Salmonella*-specific DNA hybridization probe," *APMIS* 99: 114-120, 1991.

Widjojoatmodjo et al., "Evaluation of the Magnetic Immuno PCR Assay for Rapid Detections of *Salmonella*," *European Journal of Clinical Moicrobiology and Infectious Disease* 10(11): 935-938, 1991.

Halbrook et al., "Rapid detection of salmonella in foods--a convenient two-day procedure," *Letters in Applied Microbiology* 8: 139-142, 1989.

Krysinski and Heimsch, "Use of Enzyme-Labeled Antibodies to Detect Salmonella in Foods," *Applied and Environmental Microbiology* 33(4): 947-954, 1977.

Minnich et al., "Enzyme Immunoassay for Detection of Salmonellae in Foods," *Applied and Environmental Microbiology* 43(4): 877-883, 1982.

Anderson and Hartman, "Direct Immunoassay for Detection of Salmonellae in Foods and Feeds," *Applied and Environmental Microbiology* 49(5): 1124-1127, 1985.

Cherrington and Huis in't Veld, "Comparison of classical isolation protocols with a 24 h screen to detect viable salmonellas in faeces," *Journal of Applied Bacteriology* 75: 65-68, 1993.

Feldsine et al., "Polyclonal Enzyme Immunoassay Method for Detection of Motile and Non-Motile Salmonella in Foods: Collaborative Study," *Journal of AOAC International* 75(6): 1032-1044, 1992.

Ibrahim et al., "Immobilization of Microorganisms for Detection by Solid-Phase Immunoassays," *Journal of Clinical Microbiology* 22(3): 361-365, 1985.

Ibrahim and Lyons, "Detection of Salmonellae in Foods with an Enzyme Immunometric Assay," *Journal of Food Protection* 50(1): 59-61, 1987.

Mattingly, "An Enzyme Immunoassay for the Detection of All Salmonella Using a Combination of a Myeloma Protein and a Hybridoma Antibody," *Journal of Immunological Methods* 73: 147-156, 1984.

Smith et al., "M467: A Murine IgA Myeloma Protein That Binds a Bacterial Protein. I. Recognition of Common Antigenic Determinants on Salmonella Flagellins," *Journal of Immunology* 123(4): 1715-1720, 1979.

Brooks et al., "Experimental enzyme-linked amperometric immunosensors for the detection of salmonellas in foods," *Journal of Applied Bacteriology* 73: 189-196, 1992.

Robison et al., "Enzyme Immunoassay in Which a Myeloma Protein Is Used for Detection of Salmonellae," *Applied and Environmental Microbiology* 45(6): 1816-1821, 1983.

Lee et al., "Enzyme-Linked Immunosorbent Assay for Salmonella typhimurium in Food: Feasibility of 1-Day Salmonella Detection," *Applied and Environmental Microbiology* 56(6): 1541-1546, 1990.

Flint and Hartley, "Evaluation of the TECRA immunocapture ELISA for the detection of Salmonella typhimurium in foods," *Letters in Applied Microbiology* 17: 4-6, 1993.

Isomaki et al., "Serological Diagnosis of Salmonella Infections By Enzyme Immunoassay," *Lancet* (Jun. 24): 1411-1414, 1989.

D'Aoust and Sewell, "Detection of Salmonella by the Enzyme Immunoassay (EIA) Technique," *Journal of Food Science* 51(2): 484-488, 1986.

Emswiler-Rose et al., "An Enzyme Immunoassay Technique for Detection of Salmonellae in Meat and Poultry Products," *Journal of Food Science* 49: 1018-1020, 1984.

Mattingly and Gehle, "An Improved Enzyme Immunoassay for the Detection of Salmonella," *Journal of Food Science* 49: 807-809, 1984.

Todd et al., "Development of an enzyme-linked antibody hydrophobic grid membrane filter method for the detection of Salmonella in foods," *Food Microbiology* 10: 87-99, 1993.

Todd et al., "Evaluation of Salmonella antisera for an optimum enzyme-linked antibody detection of Salmonella using hydrophobic grid membrane filters," *Food Microbiology* 8: 311-324, 1991.

van Zijderveld, "Comparison of Four Different Enzyme-Linked Immunosorbent Assays for Serological Diagnosis of Salmonella enteritidis Infections in Experimentally Infected Chickens," *Journal of Clinical Microbiology* 30(10): 2560-2566, 1992.

Araj and das Chugh, "Detection of Salmonella spp. in Clinical Specimens by Capture Enzyme-Linked Immunosorbent Assay," *Journal of Clinical Microbiology* 25: 2150-2153, 1987.

Baay and Huis in't Veld, "Alternative antigens reduce cross-reactions in an ELISA for the detection of Salmonella enteritidis in poultry," *Journal of Applied Bacteriology* 74: 243-247, 1993.

Clark et al., "Detection of Salmonella in Foods using a Novel Coloured Latex Test," *Food and Agricultural Immunology* 1: 3-9, 1989.

Benge, "Detection of Salmonella Species in Faeces by Latex Agglutination in Enrichment Broth," *European Journal Clinical Microbiol. Infect. Dis.* 8(4): 294-298, 1989.

Dadrast et al., "Egg yolk antibody detection in identification of salmonella infected poultry," *Veterinary Record* 126: 219, 1990.

Lane et al., "Detection of Food Pathogens Using the Phage Rapid Identification Assay System," *Vitek Systems Inc. and Bactomatic Inc.*

Van Poucke, "Salmonella-TEK, a Rapid Screening Method for Salmonella Species in Food," *Applied and Environmental Microbiology* 56(4): 924-927, 1990.

Lambiri et al., "Comparison of the TECRA Salmonella Immunoassay with the conventional culture method," Letters in Applied Microbiology 11: 182-184, 1990.

Humbert et al., "Rapid detection of Salmonella from poultry meat products using the '1.2. Test.RTM.," Letters in Applied Microbiology 19:245-249, 1990.

Karlsson et al., "Application and Usefulness of Enzyme Immunoassay for Diagnosis of Salmonella typhimurium Infection," Scandinavian Journal of Infectious Disease 12: 41-47, 1980.

Kerr et al., "A comparison of three salmonella antigen-capture ELISAs and culture for veterinary diagnostic specimens," Journal of Applied Bacteriology 75: 164-167, 1993.

D'Aoust and Sewell, "Detection of Salmonella with the BioEnzabead.TM. Enzyme Immunoassay Technique," Journal of Food Protection 51(7): 538-541, 1988.

Nath et al., "Evaluation of Enrichment Protocols for the 1-2Test.TM. for Salmonella Detection in Naturally Contaminated Foods & Feeds," Journal of Food Protection 52(7): 498-499, 1989.

Oggel et al., "Modified 1-2 Test.TM. System as a Rapid Screening Method for Detection of Salmonella in Foods and Feeds," Journal of Food Protection 53(8): 656-658, 1990.

Flowers et al., "Visual Immunoassay for Detection of Salmonella in Foods: Collaborative Study," Journal Assoc. Off. Anal. Chem. 71(5): 973-980, 1988.

Curiale et al., "Comparison of Colorimetric Monoclonal Enzyme Immunoassay Screening Methods for Detection of Salmonella in Foods," Journal Assoc. Off. Anal. Chem. 73(1): 43-50, 1990.

Curiale et al., "Detection of Salmonellae and Other Enterobacteriaceae by Commercial Deoxyribonucleic Acid Hybridization and Enzyme Immunoassay Kits," Journal of Food Protection 53(12): 1037-1046, 1990.

Bouvet and Jeanjean, "Evaluation of Two Colored Latex Kits, the Wellcolex Colour Salmonella Test and the Wellcox Colour Shigella Test, for Serological Grouping of Salmonella and Shigella Species," Journal of Clinical Microbiology 30(8): 2184-2186, 1992.

International BioProducts, Inc., "Executive Summary: Comparison Guide For The AOAC Accepted Rapid Methods,".

International BioProducts, Inc., "Executive Summary: Comparison Guide For The Visual Immunoassay And The 1-2 Test,".

BioControl Systems, Inc., "1-2 Test.RTM.," Technical Bulletin, Dec. 1993.

Biocontrol Systems, Inc., "Assurance.RTM. EIA: Providing Unique Advantages for Rapid Bacterial Detection,".

"Unique.TM. Salmonella," TECRA.TM. Technical Bulletin (7).

Gene-Trak Systems, "Colorimetric Gene-Trak.RTM. Salmonella Assay," 1993.

Gene-Trak Systems Corporation, "Gene-Trak.RTM. Salmonella Assay: Colorimetric protocol Overview," 1993.

Dynatech Laboratories, Inc., "Q-Trol Salmonella: Salmonella Microelisa.RTM. -92 & Microelisa.RTM. -32 Detection Kit," Catalog Nos. 007-040-0460 (92 Test) & 007-040-0440 (32 Test), 1988.

Dynatech Laboratories, Inc., "Q-Trol Salmonella: Salmonella Detection Kit," Catalog No. 007-040-0410.

"1-2 Test.RTM.--Modified per Oggel: Directions for Use,".

Pro-Lab Diagnostics, "A Vision of the Future from Pro-Lab Diagnostics," Advertisement, 1994.

AMPCOR, Inc., "DipStick Salmonella Color Change Test Kit: For Detection of Salmonella sp. in fecal specimens," Advertisement.

Murex, "Wellcolex Colour Shigella: Colour latex test for the detection and species identification of Shigella," Informational Brochure, ZC51.

Murex, "Wellcolex Colour Salmonella: Colour latex test for the detection and serogroup identification of Salmonella," Informational Brochure, ZC50.

Krusell and Skovgaard, "Evaluation of a new semi-automated screening method for the detection of Salmonella in foods within 24 h," International Journal of Food Microbiology 20: 123-130, 1993.

Incstar Corporation, "Salmonella Fluoro-Kit.TM. Delayed Direct Fluorescent Antibody Test," Instruction Manual, Catalog No. 1903, 1991.

LAB M, "Microbial Confirmation Strips," Instruction Manual and Informational Brochure.

International BioProducts, Inc., "BioPro.TM. Salmonella Visual Immunoassay: A Final Action AOAC Approved Method for the Rapid Detection of Salmonella," Informational Brochure.

Organon Teknika Corporation, "Salmonella-Tek.TM. ELISA Test System: For the detection of Salmonella antigen in foods," Instruction Manual: 1993.

AMPCOR, "Dipstick Salmonella," Instruction Manual, Catalog No. 300-150, 1991.
Pro-Lab Diagnostics, "Vision Antisera," Instruction Manual, 1993.
Wellmark Diagnostics Ltd., "Wellcolex," Advertisement.
Wellmark Diagnostics Ltd., "Salmonella Agglutinating Sera For Slide and Tube Agglutination Tests," Informational Brochure.
Mansfield and Forsythe, "Immunomagnetic separation as an alternative to enrichment broth for Salmonella detection," Letters in Applied Microbiology 16: 122-125, 1993.
Torensma et al., "Monoclonal Antibodies That Detect Live Salmonellae," Applied and Environmental Microbiology 58(12): 3868-3872, 1992.
Holbrook et al., "Comparative evaluation of the Oxoid Salmonella Rapid Test with three other rapid salmonella methods," Letters in Applied Microbiology 9: 161-164, 1989.
Izat et al., "Comparison of the DNA Probe to Culture Methods for the Detection of Salmonella on Poultry Carcasses and Processing Waters," Journal of Food Protection 52(8) 564-570, 1989.
Scholl et al., "Clinical Application of Novel Sample Processing Technology for the Identification of Salmonellae by Using DNA Probes," Journal of Clinical Microbiology 28(2): 237-241, 1990.
Rose et al., "Evaluation of a Colorimetric DNA Hybridization Test for Detection of Salmonellae in Meat and Poultry Products," Journal of Food Protection 54(2): 127-130, 1991.
Gast and Beard, "Research Note: Detection of Salmonella Serogroup D-Specific Antibodies in the Yolks of Eggs Laid by Hens Infected with Salmonella enteritidis," Poultry Science 70: 1273-1276, 1991.
Lu et al., "Characterization and application of a murine monoclonal antibody that reacts specifically with the serogroup D.sub.1 Salmonella," FEMS Microbiology Letters 80: 135-140, 1991.
McRill et al., "Application of the Peroxidase-Antiperoxidase Immunoassay to the Identification of Salmonellae from Pure Culture and Animal Tissue," Journal of Clinical Microbiology 20(2): 281-284, 1984.
Mohit et al., "A Simple Single-Step Immunoimmobilisation Method For The Detection Of Salmonella In The Presence Of Large Numbers Of Other Bacteria," Journal of Medical Microbiology 8: 173-176, 1975.
Appelmek et al., "An Enzyme-Linked Immunosorbent Assay (ELISA) for the Measurement of Antibodies to Different Parts of the Gram-Negative Lipopolysaccharide Core Region," Journal of Immunological Methods 82: 199-207, 1985.
Choi et al., "Sandwich capture ELISA by a murine monoclonal antibody against a genus-specific LPS epitope for the detection of different common serotypes of salmonellas," Journal of Applied Bacteriology 72: 134-138, 1992.
Ibrahim et al., "Immunological relationships between Salmonella flagella and their potential application for salmonellae detection by immunoassay," Med. Microbiol. Immunol. 174: 87-99, 1985.
Aleixo et al., "Enzyme Immunoassay: Binding of Salmonella Antigens to Activated Microtiter Plates," Journal of Immunoassay 6(4): 391-407, 1985.
Vermunt et al., "Isolation of salmonellas by immunomagnetic separation," Journal of Applied Bacteriology 72: 112-118, 1992.
Gruenewald et al., "Identification of Salmonella Somatic and Flagellar Antigens by Modified Serological Methods," Applied and Environmental Microbiology 56(1): 24-30, 1990.
Pilantanapak et al., "Biotinylated probes for epidemiological studies of drug resistance in Salmonella krefeld," Journal of Antimicrobial Chemotherapy 25:593-603, 1990.
Qadri et al., "Sandwich Enzyme Immunoassay for Detection of Salmonella typhi," Journal of Immunoassay 11(2): 251-270, 1990.
Luk et al., "Epitope Mapping of Four Monoclonal Antibodies Recognizing the Hexose Core Domain of Salmonella Lipopolysaccharide," Journal of Biological Chemistry 266 (34): 23215-23225, 1991.
Gruenewald et al., "Identification of Salmonella Somatic and Flagellar Antigens by Modified Serological Methods," Applied and Environmental Microbiology 56(1): 24-30, 1990.
Appassakij et al., "Enzyme-Linked Immunosorbent Assay for Detection of Salmonella typhi Protein Antigen," Journal of Clinical Microbiology 25(2): 273-277, 1987.
Muthukumar and Muthukkaruppan, "Detection of porin antigen in serum for early diagnosis of mouse infections with Salmonella typhimurium," FEMS Microbiology Immunology 89: 147-154, 1992.
Luk and Lindberg, "Anti-Salmonella Lipopolysaccharide Monoclonal Antibodies:

- Characterization of Salmonella BO-, CO-, DO-, and EO-Specific Clones and Their Diagnostic Usefulness," *Journal of Clinical Microbiology* 29(11): 2424-2433, 1991.
- Luk et al., "Characterisation and application of a murine monoclonal antibody specific for the serogroup C.sub.2 Salmonella," *J. Med. Microbiol.* 26: 115-119, 1988.
- Williams and Whittemore, "Serological Response of Chickens to Salmonella thompson and Salmonella pullorum Infections," *Journal of Clinical Microbiology* 9(1): 108-114, 1979.
- Ekpo et al., "Monoclonal Antibodies to 52-Kilodalton Protein of Salmonella typhi," *Journal of Clinical Microbiology* 28(8): 1818-1821, 1990.
- Lim, "A one-step two-particle latex immunoassay for the detection of Salmonella typhi endotoxin," *Journal of Immunological Methods* 135: 257-261, 1990.
- Bettelheim and Maskill, "Investigation by Enzyme-Linked Immunosorbent Assay of Salmonella H Antigen-Antibody Reactions," *Journal of Clinical Microbiology* 21(5): 772-774, 1985.
- Lim and Choy, "A spectrophotometric method for evaluating a latex agglutination assay of Salmonella typhi lipopolysaccharide," *Journal of Immunological Methods* 115: 269-274, 1988.
- Chaicumpa et al., "Diagnosis of Typhoid Fever by Detection of Salmonella typhi Antigen in Urine," *Journal of Clinical Microbiology* 30(9): 2513-2515, 1992.
- Caldwell et al., "Somatic and Flagellar Immunofluorescence of Salmonella," *Journal of Bacteriology* 92(4): 1177-1187, 1966.
- Pai et al., "Characterization of monoclonal antibodies to the outer membrane protein (OmpD) of Salmonella typhimurium," *Canadian Journal of Microbiology* 38: 1102-1107, 1992.
- Colwell et al., "Monoclonal Antibodies to Salmonella Lipopolysaccharide: Anti-O-Polysaccharide Antibodies Protect C3H Mice Against Challenge with Virulent Salmonella typhimurium," *Journal of Immunology* 133(2): 950-957, 1984.
- Kang et al., "Evaluation of modified passive haemagglutination assay for Vi antibody estimation in Salmonella typhi infections," *Journal of Clinical Pathology* 45:740-741, 1992.
- Luk and Lindberg, "Rapid and sensitive detection of Salmonella (O:6,7) by immunomagnetic monoclonal antibody-based assay," *Journal of Immunological Methods* 137: 1-8, 1991.
- Smith et al., "Detection of Salmonella dublin mammary gland infection in carrier cows, using an enzyme-linked immunosorbent assay for antibody in milk or serum," *Am. J. Vet. Res.* 50(8): 1352-1360, 1989.
- Ibrahim et al., "Production of Potent Salmonella H Antisera by Immunization with Polymeric Flagellins," *Journal of Clinical Microbiology* 22(3): 347-351, 1985.
- Ekwall et al., "Specific Identification of Salmonella Serogroup E Antigen O3 by Immunofluorescence and Coagulation with Antiserum Elicited by a Synthetic Trisaccharide-Bovine Serum Albumin Glycoconjugate," *Journal of Clinical Microbiology* 19(5): 699-702, 1984.
- Maxwell, "Serological Diagnosis of Salmonella Infections," *Lancet* (Aug. 19): 456, 1989.
- Barclay and Scott, "Serological Relationships between Escherichia coli and Salmonella Smooth-and Rough-Mutant Lipopolysaccharides as Revealed by Enzyme-Linked Immunosorbent Assay for Human Immunoglobulin G Antiendotoxin Antibodies," *Infection and Immunity* 55(11): 2706-2714, 1987.
- Durham et al., "Stability of Immunofluorescence Reactions Produced by Polyclonal and Monoclonal Antibody Conjugates for Rabies Virus," *Journal of Clinical Microbiology* 24(2): 301-303, 1986.
- Turpin et al., "An ion-exchange based extraction method for the detection of salmonellas in soil," *Journal of Applied Bacteriology* 74: 181-190, 1993.
- Hariharan et al., "Competitive Enzyme-Linked Immunosorbent Assay for Cholera-Related Enterotoxins in Salmonella typhimurium," *Journal of Clinical Microbiology* 24(2): 298-300, 1986.
- Chart et al., "Serological response of chickens to Salmonella enteritidis infections," *Epidemiol. Infect.* 104: 63-71, 1990.
- Stanley et al., "Molecular phylogenetic typing of pandemic isolates of Salmonella enteritidis," *FEMS Microbiology Letters* 90:153-160, 1992.
- Rubin et al., "Development of a DNA Probe to Detect Salmonella typhi," *Journal of Clinical Microbiology* 22(4): 600-605, 1985.
- Gibert et al., "Distribution of insertion sequence IS200 in Salmonella and Shigella," *Journal of General Microbiology* 136: 2555-2560, 1990.
- Altwegg et al., "Ribosomal RNA Gene Restriction Patterns Provide Increased Sensitivity

for Typing Salmonella typhi Strains," Journal of Infectious Disease 160(1): 145-149, 1989.

McLeod and Barrow, "Lipopolysaccharide-specific IgG in egg yolk from two chicken flocks infected with Salmonella enteritidis," Letters in Applied Microbiology 13: 294-297, 1991.

Kay, W. W. F. "Transport of carboxylic acids," pp. 385-411. In B. P. Rosen (eds.), Bacterial Transport. Marcel Dekker, New York, 1978.

Somers et al., "Fluorocitrate Resistant Tricarboxylate Transport Mutants of Salmonella typhimurium," Mol. Gen. Genet. 181: 338-345, 1981.

Kay and Cameron, "Transport of C.sub.4 -Dicarboxylic Acids in Salmonella typhimurium," Archives of Biochemistry and Biophysics 190(1): 281-289, 1978.

Ashton et al., "Citrate transport in Salmonella typhimurium: studies with 2-fluoro-L-erythro citrate as a substrate," Canadian Journal of Biochemistry 58(10): 797-803, 1980.

Kay et al., "Tricarboxylate Transport Systems: the tct Operon in Salmonella typhimurium," Microbiology: 34-37, 1984.

Kay and Cameron, "Citrate Transport in Salmonella typhimurium," Archives of Biochemistry and Biophysics 190(1): 270-280, 1978.

Tomas and Kay, "Tricarboxylate transport in a Cit.sup.+ Escherichia coli: evidence for the role of an outer membrane protein," Canadian Journal of Microbiology 30: 916-921, 1984.

ART-UNIT: 187

PRIMARY-EXAMINER: Campbell; Eggerton A.

ATTY-AGENT-FIRM: Seed and Berry LLP

ABSTRACT:

An isolated nucleic acid molecule comprising the agfA gene of Salmonella. Methods and compositions suitable for diagnostic tests utilizing the isolated gene, and protein therefrom, to give highly specific diagnostic assays to Salmonella, and/or enteropathogenic bacteria of the family Enterobacteriaceae.

5 Claims, 26 Drawing figures

02/18/2002

Connecting via Winsock to STN

Trying 3106016892...Open

Welcome to STN International! Enter x:x

LOGINID:sssptal806jxt

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Sep 17 IMSworld Pharmaceutical Company Directory name change
to PHARMASEARCH
NEWS 3 Oct 09 Korean abstracts now included in Derwent World Patents
Index
NEWS 4 Oct 09 Number of Derwent World Patents Index updates increased
NEWS 5 Oct 15 Calculated properties now in the REGISTRY/ZREGISTRY File
NEWS 6 Oct 22 Over 1 million reactions added to CASREACT
NEWS 7 Oct 22 DGENE GETSIM has been improved
NEWS 8 Oct 29 AAASD no longer available
NEWS 9 Nov 19 New Search Capabilities USPATFULL and USPAT2
NEWS 10 Nov 19 TOXCENTER(SM) - new toxicology file now available on STN
NEWS 11 Nov 29 COPPERLIT now available on STN
NEWS 12 Nov 29 DWPI revisions to NTIS and US Provisional Numbers
NEWS 13 Nov 30 Files VETU and VETB to have open access
NEWS 14 Dec 10 WPINDEX/WPIDS/WPIX New and Revised Manual Codes for 2002
NEWS 15 Dec 10 DGENE BLAST Homology Search
NEWS 16 Dec 17 WELDASEARCH now available on STN
NEWS 17 Dec 17 STANDARDS now available on STN
NEWS 18 Dec 17 New fields for DPCI
NEWS 19 Dec 19 CAS Roles modified
NEWS 20 Dec 19 1907-1946 data and page images added to CA and Cplus
NEWS 21 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web
NEWS 22 Jan 25 Searching with the P indicator for Preparations
NEWS 23 Jan 29 FSTA has been reloaded and moves to weekly updates
NEWS 24 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update
frequency

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 16:37:07 ON 13 FEB 2002

=> file uspat biosis caplus medline

AN 2001:301838 BIOSIS
 DN PREV200100301838
 TI Protein quantification from complex protein mixtures using a proteomics methodology with single-cell resolution.
 AU Zhang, Hong-Tao; Kacharina, Janet E.; Miyashiro, Kevin; Greene, Mark I. (1); Eberwine, James
 CS (1) Departments of Pathology and Laboratory Medicine, Abramson Institute for Cancer Research, University of Pennsylvania, Philadelphia, PA, 19104-6082: greene@reo.med.upenn.edu, eberwine@mscf.med.upenn.edu USA
 SO Proceedings of the National Academy of Sciences of the United States of America, (May 8, 2001) Vol. 98, No. 10, pp. 5497-5502. print.
 ISSN: 0027-8424.
 DT Article
 LA English
 SL English
 AB We have developed an extremely sensitive technique, termed **immuno**-detection **amplified** by T7 **RNA** polymerase (IDAT) that is capable of monitoring proteins, lipids, and metabolites and their modifications at the single-cell level. A double-stranded oligonucleotide containing the T7 promoter is conjugated to an antibody (Ab), and then T7 RNA polymerase is used to amplify RNA from the double-stranded oligonucleotides coupled to the Ab in the Ab-antigen complex. By using this technique, we are able to detect the p185her2/neu receptor from the crude lysate of T6-17 cells at 10-13 dilution, which is 109-fold more sensitive than the conventional ELISA method. Single-chain Fv fragments or complementarity determining region peptides of the Ab also can be substituted for the Ab in IDAT. In a modified protocol, the oligonucleotide has been coupled to an Ab against a common epitope to create a universal detector species. With the linear **amplification** ability of T7 **RNA** polymerase, IDAT represents a significant improvement over **immuno**-PCR in terms of sensitivity and has the potential to provide a robotic platform for proteomics.

L2 ANSWER 3 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 AN 2002:4350 BIOSIS
 DN PREV200200004350
 TI A novel proteomics methodology: **Immuno**-detection **amplified** by T7 **RNA** polymerase.
 AU Kacharina, J. Estee (1); Zhang, H.; Greene, M.; Eberwine, J. (1)
 CS (1) Pharmacology, University of Pennsylvania, Philadelphia, PA USA
 SO Society for Neuroscience Abstracts, (2001) Vol. 27, No. 2, pp. 2451. print.
 Meeting Info.: 31st Annual Meeting of the Society for Neuroscience San Diego, California, USA November 10-15, 2001
 ISSN: 0190-5295.
 DT Conference
 LA English
 AB In our continuing efforts to understand the molecular underpinnings of dendritic functioning, we are expanding our studies to include the simultaneous analysis of multiple proteins in neurons. Towards this end we have utilized a novel proteomics methodology termed **immuno**-detection **amplified** by T7 **RNA** polymerase (IDAT) that can quantify the level of multiple proteins in a single cell. This methodology is a modified sandwich ELISA in which Ab-antigen interactions are detected by nucleic acid amplification. A ds-oligo containing a T7 promoter site was covalently coupled to a detector Ab or single-chain Fv fragment and served as the cDNA template. After interaction of the protein target with the detector Ab, RNA amplification was performed. The resulting RNA was labeled with Ribo-Green, and measured with a fluorescent microplate reader. IDAT has recently been used to measure the amount of p185her2/neu, a protein marker for breast cancer, in 3T3 cells and in a single rat hippocampal neuron. We are currently generating phage display peptide libraries enriched for rat hippocampal neuronal or post-synaptic density proteins to be used in conjunction with the IDAT technology. This

proteomics approach will make it possible to screen neurons for the presence and abundance of thousands of proteins and their post-translational modifications in response to neuronal modulation.

L2 ANSWER 4 OF 9 USPATFULL
AN 1999:18920 USPATFULL
TI Method for amplification and detection of RNA and DNA sequences
IN Rossi, John J., Glendora, CA, United States
PA City of Hope, Duarte, CA, United States (U.S. corporation)
PI US 5869249 19990209
AI US 1996-747489 19961112 (8)
RLI Continuation of Ser. No. US 1994-334398, filed on 3 Nov 1994, now patented, Pat. No. US 5622820 which is a continuation of Ser. No. US 1988-180740, filed on 12 Apr 1988, now abandoned which is a continuation-in-part of Ser. No. US 1988-165915, filed on 10 Mar 1988, now abandoned
DT Utility
FS Granted
EXNAM Primary Examiner: Zitomer, Stephanie W.
LREP Rothwell, Figg, Ernst & Kurz
CLMN Number of Claims: 7
ECL Exemplary Claim: 1
DRWN 17 Drawing Figure(s); 13 Drawing Page(s)
LN.CNT 763
CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A method of detecting and identifying specific human nucleic acid sequences contained in human nucleic acids in a human blood or tissue sample, comprising: (1) amplifying at least one portion of the specific nucleic acid sequence present in the sample; (2) transcribing the amplification product with an RNA polymerase to produce multiple RNA copies of each copy of specific nucleic acid sequence comprising the amplification product; and (3) identifying the transcription product.

L2 ANSWER 5 OF 9 USPATFULL
AN 1998:150654 USPATFULL
TI Hepatitis C virus epitopes
IN Reyes, Gregory R., Palo Alto, CA, United States
Kim, Jungsuh P., Palo Alto, CA, United States
Moeckli, Randolph, Redwood City, CA, United States
Simonsen, Christian C., Saratoga, CA, United States
PA Genelabs Technologies, Inc., Redwood City, CA, United States (U.S. corporation)
PI US 5843636 19981201
AI US 1995-407410 19950317 (8)
RLI Division of Ser. No. US 1991-681703, filed on 5 Apr 1991, now patented, Pat. No. US 5443965 which is a continuation-in-part of Ser. No. US 1990-594854, filed on 10 Oct 1990, now abandoned which is a continuation-in-part of Ser. No. US 1990-205611, filed on 6 Apr 1990, now abandoned
DT Utility
FS Granted
EXNAM Primary Examiner: Woodward, Michael P.
LREP Dehlinger & Associates
CLMN Number of Claims: 12
ECL Exemplary Claim: 1
DRWN 21 Drawing Figure(s); 21 Drawing Page(s)
LN.CNT 3115
CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Peptide antigens which are immunoreactive with sera from individuals infected with hepatitis C virus (HCV) are disclosed. Several of the antigens are immunologically reactive with antibodies present in individuals identified as having chronic and acute HCV infection. The antigens are useful in diagnostic methods for detecting HCV infection in humans. Also disclosed are corresponding genomic-fragment clones

containing polynucleotides encoding the open reading frame sequences for the antigenic peptides.

L2 ANSWER 6 OF 9 MEDLINE
AN 96376777 MEDLINE
DN 96376777 PubMed ID: 8782649
TI Detection of N-acetylgalactosaminyltransferase mRNA which determines expression of Sda blood group carbohydrate structure in human gastrointestinal mucosa and cancer.
AU Dohi T; Yuyama Y; Natori Y; Smith P L; Lowe J B; Oshima M
CS Biochemistry and Nutrition Division, Research Institute, International Medical Center of Japan, Tokyo.
SO INTERNATIONAL JOURNAL OF CANCER, (1996 Sep 4) 67 (5) 626-31.
Journal code: GQU; 0042124. ISSN: 0020-7136.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
OS GENBANK-S83275
EM 199610
ED Entered STN: 19961106
Last Updated on STN: 19980206
Entered Medline: 19961024
AB The Sda blood group carbohydrate structure, GalNAcbeta1-4[NeuAcalpha2-3]Galbeta1-4GlcNAc-R, is expressed on glycolipid and glycoprotein in human gastrointestinal mucosa. The expression of the Sda determinant dramatically decreases in cancer tissue. The activity of the beta1,4N-acetylgalactosaminyltransferase (Sda-GalNAcT), which transfers GalNAc to NeuAcalpha2-3Galbeta1-4Glc(NAc)-R, correlates with the expression of the Sda immuno-epitope. From the total RNA fraction of human gastric mucosa, we have **amplified** a cDNA segment by reverse-transcription-polymerase-chain reaction (RT-PCR), using primers designed according to the cDNA sequence of a murine beta1,4GalNAcT which synthesizes the Sda determinant. An RT-PCR product of 390 bp shared 85% nucleotide identity with the murine Sda-related beta1,4GalNAcT. This RT-PCR product hybridized to a transcript in mRNA prepared from human gastric mucosa. In RT-PCR using specific primers to this PCR product, Sda-GalNAcT mRNA was detected in all samples of normal stomach and small intestine examined and the majority of normal colonic specimens. Six out of nine cases of gastric cancer, and 9 out of 13 cases of colonic cancer failed to produce the target DNA. These results correlate with the beta1,4GalNAcT activity measured in the same samples. In conclusion, a segment of the cDNA for beta1,4GalNAcT which determines expression of the Sda carbohydrate structure was obtained, and reduced transcription of this beta 1,4GalNAcT resulted in the disappearance of the Sda epitope in gastrointestinal cancer.

L2 ANSWER 7 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE 2
AN 1992:326726 BIOSIS
DN BA94:28567
TI ISOLATION AND AMPLIFICATION OF HUMAN IGE FD ENCODING MRNA FROM HUMAN PERIPHERAL BLOOD LYMPHOCYTES.
AU WALKER M R; BEVAN L J; DANIELS J; ROTTIER M M A; RAPLEY R; ROBERTS A M
CS MOLECULAR BIOLOGY RES. GROUP, UNIVERSITY DEP. CLINICAL BIOCHEMISTRY, WOLFSON RES. LAB., QUEEN ELIZABETH MEDICAL CENTRE, EDGBASTON, BIRMINGHAM B15 2TH, UK.
SO J IMMUNOL METHODS, (1992) 149 (1), 77-85.
CODEN: JIMMBG. ISSN: 0022-1759.
FS BA; OLD
LA English
AB In order to establish the feasibility of applying recombinatorial library technologies to investigate human in vivo IgE responses, and as a pre-requisite of recombinatorial library construction, we have attempted to determine workable peripheral blood sample volumes required for

isolation of mRNA for polymerase chain reaction (PCR) amplification of human IgE Fd encoding sequences. Cells secreting chimeric human IgE monoclonal antibody specific for the hapten NIP were used to establish the conditions for specific amplification of C.epsilon.1 domain and Fd encoding sequences, as determined by Southern hybridisation. Amplification of C.epsilon.1 domain sequences could be achieved using as few as ten cultured cells as the source of RNA. Specific IgE+ B cell enrichment using **immuno**-magnetic particles prior to **RNA** extraction was, however, required to obtain **amplification** of IgE C.epsilon.1 and Fd fragments from lymphocytes prepared from 40 ml human peripheral blood. IgG1+ B cell enrichment from similar samples was not required for detectable amplification of human C.gamma.1 cDNA sequences. However, this procedure improved amplification efficiency. Optimisation of methods to separate specific B cell populations, or specific RNA/cDNA sequences, will facilitate in vitro generation of human IgE Fab fragments from peripheral blood.

L2 ANSWER 8 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 AN 1983:291438 BIOSIS
 DN BA76:48930
 TI CONSTRUCTION OF A MODULAR DI HYDRO FOLATE REDUCTASE COMPLEMENTARY DNA GENE ANALYSIS OF SIGNALS UTILIZED FOR EFFICIENT EXPRESSION.
 AU KAUFMAN R J; SHARP P A
 CS DEP. BIOL., MASS. INST. TECHNOL., CAMBRIDGE, MASS. 02139.
 SO MOL CELL BIOL, (1982) 2 (11), 1304-1319.
 CODEN: MCEBD4. ISSN: 0270-7306.
 FS BA; OLD
 LA English
 AB Dihydrofolate reductase (DHFR) modular genes were constructed with segments containing the adenovirus major late promoter, a 3' splice site from a variable region Ig gene, a DHFR c[complementary]DNA, and portions of the SV40 genome. DNA-mediated transfer of these genes transformed Chinese hamster ovary [CHO] DHFR- cells to the DHFR+ phenotype. Transformants contained one to several copies of the transfected DNA integrated into the host genome. Clones subjected to growth in increasing concentrations of methotrexate eventually gave rise to lines containing several hundred copies of the transforming DNA. Analysis of the DHFR mRNA produced in amplified lines indicated the following. All clones utilize the adenovirus major late promoter for transcription initiation. A hybrid intron formed by the 5' splice site of the adenovirus major late leader and a 3' splice site from a variable-region Ig gene is properly excised. The mRNA is not efficiently polyadenylated at sequences in the 3' end of the DHFR cDNA but rather uses polyadenylation signals downstream from the DHFR cDNA. Three independent clones produce a DHFR mRNA containing SV40 or pBR322 and SV40 sequences, and the RNA is polyadenylated at the SV40 late polyadenylation site. Another clone has recombined into cellular DNA and apparently uses a cellular sequence for polyadenylation. Introduction of a segment containing the SV40 early polyadenylation signal into the 3' end of the DHFR cDNA gene generated a recombinant capable of transforming cells to the DHFR+ phenotype with at least a 10-fold increase in efficiency, demonstrating the necessity for an efficient polyadenylation signal. Attachment of a DNA segment containing the transcription enhancer (72-base pair repeat) of SV40 further increased the biological activity of the modular DHFR gene 50- to 100-fold.

L2 ANSWER 9 OF 9 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 AN 1981:176634 BIOSIS
 DN BA71:46626
 TI A HIGH PRODUCTION RATE OF TRANSLATABLE **IMMUNO** GLOBULIN G MESSENGER **RNA** ACCOUNTS FOR THE **AMPLIFIED** SYNTHESIS OF **IMMUNO** GLOBULIN G IN MYELOMA CELLS.
 AU WALLACH M; LASKOV R
 CS THE HUBERT H. HUMPHREY CENTER FOR EXPERIMENTAL MEDICINE AND CANCER RESEARCH, THE HEBREW UNIVERSITY-HADASSAH MEDICAL SCHOOL, JERUSALEM,

ISRAEL.

SO EUR J BIOCHEM, (1980) 110 (2), 545-554.
CODEN: EJBACI. ISSN: 0014-2956.

FS BA; OLD

LA English

AB The accumulation of Ig mRNA in myeloma cells was studied. Specific mRNA for L and H chains of IgG were isolated from the murine MPC-11 myeloma tumor cells by immune precipitation of polysomes which synthesize these chains. The immune-precipitated polysomes were enriched 10- to 30-fold in the .gamma. and .vkappa. mRNA sequences, respectively. In the wheat germ cell-free system the .vkappa. mRNA preparation was translated mainly into 3 polypeptides of MW 25,000, 18,000 and 15,000. Immune precipitation of polysomes was also used to characterize 3 variant clones of MPC-11 myeloma. Little if any .gamma.-chain polysomes are present in the L-chain producer and non-producer clones, while a substantial amount of .vkappa.-chain polysomes was present in the non-producer clone. This may be due to the presence in the non-producer cells of the constant region .vkappa.-chain fragment. In order to determine the relative synthesis rate of .vkappa. and .gamma. mRNA, pulse-labeled polysomes were immune precipitated using antibodies to .vkappa. and .gamma. chains. .vkappa. and .gamma. mRNA molecules are produced at a very high relative rate each accounting for 10-15% of the total labeled mRNA after 1 h of labeling. These values are higher than the steady-state pool size of .vkappa. and .gamma. mRNA, which was 5-6%, and indicates that the half-life of these molecules is not unusually high. The amplified synthesis of Ig chains in myeloma cells is mainly due to a high rate of production of translatable .vkappa. and .gamma. mRNA.

=>

L2 ANSWER 4 OF 9 USPATFULL

SUMM **Amplification** of an **RNA** template is particularly advantageous for detecting retroviruses, such as human **immuno**-deficiency virus (HIV or human T-lymphotropic virus HTLV III/LAV). HIV has been shown to be the etiological agent of acquired immune. . .

L2 ANSWER 5 OF 9 USPATFULL

DETD . . . as 409-1-1(c-a), do yield HCV-specific immunopositive signals with selected samples. Accordingly, the two methods presented in the present invention, (i) **immuno**-screening of cDNA libraries generated directly from sera-derived **RNA**, and (ii) immunoscreening of **amplified**-fragment libraries, can both be seen to be effective methods of identifying cDNA sequences encoding viral antigens. Further, confirmation of the. . .